

## SUMMARY OF PRODUCT CHARACTERISTICS

### 1. Name of the medicinal product

SANFUR – 750 (Cefuroxime for Injection USP 750mg)

### 2. Qualitative and quantitative composition

Each vial contains:

Sterile Cefuroxime Sodium USP equivalent to anhydrous Cefuroxime 750mg

### 3. Pharmaceutical form

Dry powder for injection,

**Appearance of powder:** Sterile, white or faintly yellow powder, distributed in sealed container and which when shaken with the prescribed volume of sterile liquid, rapidly form clear and practically particle-free solution

## 4. CLINICAL PARTICULARS

### Therapeutic indications

Cefuroxime for Injection is indicated for the treatment of the infections listed below in adults and children, including neonates (from birth).

- Community acquired pneumonia
- Acute exacerbations of chronic bronchitis
- Complicated urinary tract infections, including pyelonephritis
- Soft-tissue infections: cellulitis, erysipelas and wound infections
- Intra-abdominal infections
- Prophylaxis against infection in gastrointestinal (including oesophageal), orthopaedic, cardiovascular, and gynecological surgery (including caesarean section)

In the treatment and prevention of infections in which it is very likely that anaerobic organisms will be encountered, cefuroxime should be administered with additional appropriate antibacterial agents.

Consideration should be given to official guidance on the appropriate use of antibacterial agents.

## Posology and method of administration

### Posology

Table 1. Adults and children  $\geq 40$  kg

| Indication  | Dosage   |
|---|--|
| Community acquired pneumonia and acute exacerbations of chronic bronchitis  | 750 mg every 8 hours (intravenously or intramuscularly)  |
| Soft-tissue infections: cellulitis, erysipelas and wound infections.  |  |
| Intra-abdominal infections  |  |
| Complicated urinary tract infections, including pyelonephritis  | 1.5 g every 8 hours (intravenously or intramuscularly)   |
| Severe infections   | 750 mg every 6 hours (intravenously)<br>1.5 g every 8 hours (intravenously)  |
| Surgical prophylaxis for gastrointestinal, gynecological surgery (including caesarean section) and orthopaedic operations | 1.5 g with the induction of anaesthesia. This may be supplemented with two 750 mg doses (intramuscularly) after 8 hours and 16 hours |
| Surgical prophylaxis for cardiovascular and oesophageal operations  | 1.5 g with induction of anaesthesia followed by 750 mg (intramuscularly) every 8 hours for a further 24 hours                        |

Table 2. Children < 40 kg

|   | Infants and toddlers > 3 weeks and children < 40 kg  | Infants (birth to 3 weeks)  |
|---|--|---|
| Community acquired pneumonia  | 30 to 100 mg/kg/day (intravenously) given as 3 or 4 divided doses; a dose of 60 mg/kg/day is appropriate for most infections | 30 to 100 mg/kg/day (intravenously) given as 2 or 3 divided doses |
| Complicated urinary tract infections, including pyelonephritis      |  |   |
| Soft-tissue infections: cellulitis, erysipelas and wound infections |  |   |
| Intra-abdominal infections  |  |   |

### Renal impairment

Cefuroxime is primarily excreted by the kidneys. Therefore, as with all such antibiotics, in patients with markedly impaired renal function it is recommended that the dosage of Cefuroxime for Injection should be reduced to compensate for its slower excretion.

Table 3. Recommended doses for Cefuroxime for Injection in renal impairment

| Creatinine clearance  | T <sub>1/2</sub> (hrs)      | Dose (mg)  |
|---|-----------------------------|--|
| > 20 mL/min/1.73 m <sup>2</sup>   | 1.7-2.6                     | It is not necessary to reduce the standard dose (750 mg to 1.5 g three times daily).   |
| 10-20 mL/min/1.73 m <sup>2</sup>  | 4.3-6.5                     | 750 mg twice daily   |
| < 10 mL/min/1.73 m <sup>2</sup>   | 14.8-22.3                   | 750 mg once daily  |
| Patients on haemodialysis   | 3.75                        | A further 750 mg dose should be given intravenously or intramuscularly at the end of each dialysis; in addition to parenteral use, Cefuroxime sodium can be incorporated into the peritoneal dialysis fluid (usually 250 mg for every 2 litres of dialysis fluid). |
| Patients in renal failure on Continuous arteriovenous haemodialysis (CAVH) or high-flux haemofiltration (HF) in intensive therapy units | 7.9-12.6 (CAVH)<br>1.6 (HF) | 750 mg twice daily; for low-flux haemofiltration follow the dosage recommended under impaired renal function.  |

### Hepatic impairment

Cefuroxime is primarily eliminated by the kidney. In patients with hepatic dysfunction this is not expected to affect the pharmacokinetics of cefuroxime.

### Method of administration

Cefuroxime should be administered by intravenous injection over a period of 3 to 5 minutes directly into a vein or via a drip tube or infusion over 30 to 60 minutes, or by deep intramuscular injection.

Intramuscular injections should be injected well within the bulk of a relatively large muscle and not more than 750 mg should be injected at one site. For doses greater than 1.5 g intravenous administration should be used. For instructions on reconstitution of the medicinal product

before administration.

For instructions on preparation of the medicinal product before administration.

### **Contraindications**

Hypersensitivity to cefuroxime

Patients with known hypersensitivity to cephalosporin antibiotics

History of severe hypersensitivity (e.g., anaphylactic reaction) to any other type of beta-lactam antibacterial agent (penicillins, monobactams and carbapenems)

## **Special warnings and precautions for use**

### Hypersensitivity reactions

As with all beta-lactam antibacterial agents, serious and occasionally fatal hypersensitivity reactions have been reported. In case of severe hypersensitivity reactions, treatment with cefuroxime must be discontinued immediately and adequate emergency measures must be initiated.

Before beginning treatment, it should be established whether the patient has a history of severe hypersensitivity reactions to cefuroxime, to other cephalosporins or to any other type of beta-lactam agent. Caution should be used if cefuroxime is given to patients with a history of non-severe hypersensitivity to other beta-lactam agents.

### Concurrent treatment with potent diuretics or aminoglycosides

Cephalosporin antibiotics at high dosage should be given with caution to patients receiving concurrent treatment with potent diuretics such as furosemide or aminoglycosides. Renal impairment has been reported during use of these combinations. Renal function should be monitored in the elderly and those with known pre-existing renal impairment.

### Overgrowth of non-susceptible microorganisms

Use of cefuroxime may result in the overgrowth of *Candida*. Prolonged use may also result in the overgrowth of other non-susceptible microorganisms (e.g., enterococci and *Clostridium difficile*), which may require interruption of treatment.

Antibacterial agent-associated pseudomembranous colitis has been reported with use of cefuroxime and may range in severity from mild to life threatening. This diagnosis should be considered in patients with diarrhoea during or subsequent to the administration of cefuroxime. Discontinuation of therapy with cefuroxime and the administration of specific treatment for *Clostridium difficile* should be considered. Medicinal products that inhibit peristalsis should not be given.

### Intra-abdominal infections

Due to its spectrum of activity, cefuroxime is not suitable for the treatment of infections caused by Gram-negative non-fermenting bacteria.

### Interference with diagnostic tests

The development of a positive Coomb's Test associated with the use of cefuroxime may interfere with cross matching of blood.

Slight interference with copper reduction methods (Benedict's, Fehling's, Clinitest) may be observed. However, this should not lead to false-positive results, as may be experienced with some other cephalosporins.

As a false negative result may occur in the ferricyanide test, it is recommended that either the glucose oxidase or hexokinase methods are used to determine blood/plasma glucose levels in patients receiving cefuroxime sodium.

### **Interactions with other Medicinal Products and other forms of Interaction**

Cefuroxime may affect the gut flora, leading to lower oestrogen reabsorption and reduced efficacy of combined oral contraceptives.

Cefuroxime is excreted by glomerular filtration and tubular secretion. Concomitant use of probenecid is not recommended. Concurrent administration of probenecid prolongs the excretion of the antibiotic and produces an elevated peak serum level.

### **Potential nephrotoxic drugs and loop diuretics**

High-dosage treatments with cephalosporins should be carried out with caution on patients who are taking strong-acting diuretics (such as furosemide) or potential nephrotoxic preparations (such as aminoglycoside antibiotics), since impairment of renal function through such combinations cannot be ruled out.

## **Fertility, Pregnancy and Lactation**

### Fertility

There are no data on the effects of cefuroxime sodium on fertility in humans. Reproductive studies in animals have shown no effects on fertility.

### Pregnancy

There are limited amounts of data from the use of cefuroxime in pregnant women. Studies in animals have shown no reproductive toxicity. Cefuroxime should be prescribed to pregnant women only if the benefit outweighs the risk.

Cefuroxime has been shown to cross the placenta and attain therapeutic levels in amniotic fluid and cord blood after intramuscular or intravenous dose to the mother.

### Lactation

Cefuroxime is excreted in human milk in small quantities. Adverse reactions at therapeutic doses are not expected, although a risk of diarrhoea and fungus infection of the mucous membranes cannot be excluded. A decision must be made whether to discontinue breast-feeding or to discontinue/abstain from cefuroxime therapy taking into account the benefit of breast feeding for the child and the benefit of therapy for the woman.

## **Effects on Ability to Drive and Use Machines**

No studies on the effects of cefuroxime on the ability to drive and use machines have been performed. However, based on known adverse reactions, cefuroxime is unlikely to have an effect on the ability to drive and use machines.

## **4.8 Undesirable Effects**

The most common adverse reactions are neutropenia, eosinophilia, transient rise in liver enzymes or bilirubin, particularly in patients with pre-existing liver disease, but there is no evidence of harm to the liver and injection site reactions.

The frequency categories assigned to the adverse reactions below are estimates, as for most reactions suitable data for calculating incidence are not available. In addition, the incidence of adverse reactions associated with cefuroxime sodium may vary according to the indication.

Data from clinical trials were used to determine the frequency of very common to rare adverse reactions. The frequencies assigned to all other adverse reactions (i.e., those occurring at <1/10,000) were mainly determined using post-marketing data, and refer to a reporting rate rather than a true frequency.

Treatment related adverse reactions, all grades, are listed below by MedDRA body system organ class, frequency and grade of severity. The following convention has been utilized for the classification of frequency: very common  $\geq 1/10$ ; common  $\geq 1/100$  to  $< 1/10$ ; uncommon  $\geq 1/1,000$  to  $< 1/100$ ; rare  $\geq 1/10,000$  to  $< 1/1,000$ ; very rare  $< 1/10,000$  and not known (cannot be estimated from the available data).

| System organ class                   | Common   | Uncommon                          | Not known   |
|--------------------------------------|--|-----------------------------------|---|
| Infections and infestations          |  |                                   | Candida overgrowth, overgrowth of Clostridium difficile               |
| Blood and lymphatic system disorders | neutropenia, eosinophilia, decreased haemoglobin concentration | leukopenia, positive Coomb's test | thrombocytopenia, haemolytic anaemia                                  |
| Immune disorders system              |  |                                   | drug fever, interstitial nephritis, anaphylaxis, cutaneous vasculitis |
| Gastrointestinal disorders           |  | gastrointestinal disturbance      | pseudomembranous colitis  |
| Hepatobiliary disorders              | transient rise in liver  | transient rise in                 |   |



|  | enzymes  | bilirubin                         |  |
|--|--|-----------------------------------|--|
| Skin and subcutaneous tissue disorders               |  | skin rash, urticaria and pruritus | erythema multiforme, toxic epidermal necrolysis and Stevens-Johnson syndrome, angioneurotic oedema   |
| Renal and urinary disorders                          |  |                                   | elevations in serum creatinine, elevations in blood urea nitrogen and decreased creatinine clearance |
| General disorders and administration site conditions | injection site reactions which may include pain and thrombophlebitis |                                   |  |

#### Description of selected adverse reactions

Cephalosporins as a class tend to be absorbed onto the surface of red cell membranes and react with antibodies directed against the drug to produce a positive Coomb's test (which can interfere with cross matching of blood) and very rarely haemolytic anaemia.

Transient rises in serum liver enzymes or bilirubin have been observed which are usually reversible.

Pain at the intramuscular injection site is more likely at higher doses. However, it is unlikely to be a cause for discontinuation of treatment.

#### *Paediatric population*

The safety profile for cefuroxime sodium in children is consistent with the profile in adults.

### Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorization of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product.

#### **4.9. Overdose**

Overdose can lead to neurological sequelae including encephalopathy, convulsions and coma. Symptoms of overdose can occur if the dose is not reduced appropriately in patients with renal impairment.

Serum levels of cefuroxime can be reduced by hemodialysis or peritoneal dialysis.

## **5. PHARMACOLOGICAL PROPERTIES**

### **Pharmacodynamic properties**

Pharmacotherapeutic group: Antibacterials for systemic use, second-generation cephalosporins, ATC code: J01DC02

### Mechanism of action

Cefuroxime inhibits bacterial cell wall synthesis following attachment to penicillin binding proteins (PBPs). This results in the interruption of cell wall (peptidoglycan) biosynthesis, which leads to bacterial cell lysis and death.

### Mechanism of resistance

Bacterial resistance to cefuroxime may be due to one or more of the following mechanisms:

- hydrolysis by beta-lactamases including (but not limited to) extended-spectrum beta-lactamases (ESBLs), and Amp-C enzymes, that may be induced or stably derepressed in certain aerobic Gram-negative bacterial species;
- reduced affinity of penicillin-binding proteins for cefuroxime;

- outer membrane impermeability, which restricts access of cefuroxime to penicillin binding proteins in Gram-negative bacteria;
- bacterial efflux pumps

Organisms that have acquired resistance to other injectable cephalosporins are expected to be resistant to cefuroxime. Depending on the mechanism of resistance, organisms with acquired resistance to penicillins may demonstrate reduced susceptibility or resistance to cefuroxime.

#### Cefuroxime sodium breakpoints

Minimum inhibitory concentration (MIC) breakpoints established by the European Committee on Antimicrobial Susceptibility Testing (EUCAST) are as follows:

| Microorganism                                | Breakpoints (mg/L) |                   |
|--|--------------------|-------------------|
|  | <u>Susceptible</u> | <u>Resistant</u>  |
| <i>Enterobacteriaceae</i> <sup>1</sup>       | ≤8 <sup>2</sup>    | >8                |
| <i>Staphylococcus</i> spp.                   | Note <sup>3</sup>  | Note <sup>3</sup> |
| <i>Streptococcus</i> A, B, C and G           | Note <sup>4</sup>  | Note <sup>4</sup> |
| <i>Streptococcus pneumoniae</i>              | ≤0.5               | >1                |
| <i>Streptococcus</i> (other)                 | ≤0.5               | >0.5              |
| <i>Haemophilus influenzae</i>                | ≤1                 | >2                |
| <i>Moraxella catarrhalis</i>                 | ≤4                 | >8                |
| Non-species related breakpoints <sup>1</sup> | ≤4 <sup>5</sup>    | >8 <sup>5</sup>   |

<sup>1</sup> The cephalosporin breakpoints for *Enterobacteriaceae* will detect all clinically important resistance mechanisms (including ESBL and plasmid mediated AmpC). Some strains that produce beta-lactamases are susceptible or intermediate to 3<sup>rd</sup> or 4<sup>th</sup> generation cephalosporins with these breakpoints and should be reported as found, i.e., the presence or absence of an ESBL does not in itself influence the categorization of susceptibility. In many areas, ESBL detection

and characterization is recommended or mandatory for infection control purposes.

<sup>2</sup> Breakpoint relates to a dosage of 1.5 g × 3 and to *E. coli*, *P. mirabilis* and *Klebsiella* spp. only

<sup>3</sup> Susceptibility of staphylococci to cephalosporins is inferred from the methicillin susceptibility except for ceftazidime and cefixime and ceftibuten, which do not have breakpoints and should not be used for staphylococcal infections.

<sup>4</sup> The susceptibility of streptococcus groups A, B, C and G to cephalosporins is inferred from the benzylpenicillin susceptibility.

<sup>5</sup> Breakpoints apply to daily intravenous dose of 750 mg × 3 and a high dose of at least 1.5 g × 3.

### Microbiological susceptibility

The prevalence of acquired resistance may vary geographically and with time for selected species and local information on resistance is desirable, particularly when treating severe infections. As necessary, expert advice should be sought when the local prevalence of resistance is known and the utility of the agent in at least some types of infections is questionable.

Cefuroxime is usually active against the following microorganisms *in vitro*.

#### **Commonly susceptible species**

##### Gram-positive aerobes:

*Staphylococcus aureus* (methicillin-susceptible) §

*Streptococcus pyogenes*

*Streptococcus agalactiae*

##### Gram-negative aerobes:

*Haemophilus parainfluenzae*

*Moraxella catarrhalis*

#### **Microorganisms for which acquired resistance may be a problem**

##### Gram-positive aerobes:

*Streptococcus pneumoniae*

*Streptococcus mitis* (viridans group)

Gram-negative aerobes:

*Citrobacter* spp. not including *C. freundii*

*Enterobacter* spp. not including *E. aerogenes* and *E. cloacae*

*Escherichia coli*

*Haemophilus influenzae*

*Klebsiella pneumoniae*

*Proteus mirabilis*

*Proteus* spp. not including *P. penneri* and *P. Vulgaris*

*Providencia* spp.

*Salmonella* spp.

Gram-positive

anaerobes:

*Peptostreptococcus* spp.

*Propionibacterium* spp.

Gram-negative anaerobes:

*Fusobacterium* spp.

*Bacteroides* spp.

**Inherently resistant microorganisms**

Gram-positive aerobes:

*Enterococcus faecalis*

*Enterococcus faecium*

Gram-negative aerobes:

*Acinetobacter* spp.

*Burkholderia cepacia*

*Campylobacter* spp.

*Citrobacter freundii*

*Enterobacter aerogenes*

*Enterobacter cloacae*

|   |
|---|
| <i>Morganella morganii</i><br><i>Proteus penneri</i><br><i>Proteus vulgaris</i><br><i>Pseudomonas aeruginosa</i><br><i>Serratia marcescens</i><br><i>Stenotrophomonas maltophilia</i> |
| <u>Gram-positive anaerobes:</u><br><br><i>Clostridium difficile</i>   |
| <u>Gram-negative anaerobes:</u><br><br><i>Bacteroides fragilis</i>  |
| <u>Others:</u><br><br><i>Chlamydia</i> spp.<br><i>Mycoplasma</i> spp.<br><i>Legionella</i> spp.   |

\$ All methicillin-resistant *S. aureus* are resistant to cefuroxime.

*In vitro* the activities of cefuroxime sodium and aminoglycoside antibiotics in combination have been shown to be at least additive with occasional evidence of synergy.

### **Pharmacokinetic properties**

#### Absorption

After intramuscular (IM) injection of cefuroxime to normal volunteers, the mean peak serum concentrations ranged from 27 to 35 µg/mL for a 750 mg dose and from 33 to 40 µg/mL for a 1000 mg dose, and were achieved within 30 to 60 minutes after administration. Following intravenous (IV) doses of 750 and 1500 mg, serum concentrations were approximately 50 and 100 µg/mL, respectively, at 15 minutes.

AUC and C<sub>max</sub> appear to increase linearly with increase in dose over the single dose range of 250 to 1000 mg following IM and IV administration. There was no evidence of accumulation of cefuroxime in the serum from normal volunteers following repeat intravenous administration of 1500 mg doses every 8 hours.

### Distribution

Protein binding has been stated as 33 to 50%, depending on the methodology used. The average volume of distribution ranges from 9.3 to 15.8 L/1.73 m<sup>2</sup> following IM or IV administration over the dosage range of 250 to 1000 mg. Concentrations of cefuroxime in excess of the minimum inhibitory levels for common pathogens can be achieved in the tonsilla, sinus tissues, bronchial mucosa, bone, pleural fluid, joint fluid, synovial fluid, interstitial fluid, bile, sputum and aqueous humour. Cefuroxime passes the blood-brain barrier when the meninges are inflamed.

### Biotransformation

Cefuroxime is not metabolised

### Elimination

Cefuroxime is excreted by glomerular filtration and tubular secretion. The serum half-life after either intramuscular or intravenous injection is approximately 70 minutes. There is an almost complete recovery (85 to 90%) of unchanged cefuroxime in urine within 24 hours of administration. The majority of the cefuroxime is excreted within the first 6 hours. The average renal clearance ranges from 114 to 170 mL/min/1.73 m<sup>2</sup> following IM or IV administration over the dosage range of 250 to 1000 mg.

### *Special patient populations*

#### Gender

No differences in the pharmacokinetics of cefuroxime were observed between males and females following a single IV bolus injection of 1000 mg of cefuroxime as the sodium salt.

#### Elderly

Following IM or IV administration, the absorption, distribution and excretion of cefuroxime in elderly patients are similar to younger patients with equivalent renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in cefuroxime dose selection, and it may be useful to monitor renal function.

#### Paediatrics

The serum half-life of cefuroxime has been shown to be substantially prolonged in neonates according to gestational age. However, in older infants (aged >3 weeks) and in children, the serum half-life of 60 to 90 minutes is similar to that observed in adults.

#### Renal impairment

Cefuroxime is primarily excreted by the kidneys. As with all such antibiotics, in patients with

markedly impaired renal function (i.e.,  $Cl_{cr} < 20$  mL/minute) it is recommended that the dosage of cefuroxime should be reduced to compensate for its slower excretion. Cefuroxime is effectively removed by haemodialysis and peritoneal dialysis.

#### Hepatic impairment

Since cefuroxime is primarily eliminated by the kidney, hepatic dysfunction is not expected to have an effect on the pharmacokinetics of cefuroxime.

### **5.3. Preclinical Safety Data**

Non-clinical data reveal no special hazard for humans based on conventional studies of safety pharmacology, repeated dose toxicity, genotoxicity and toxicity to reproduction and development. No carcinogenicity studies have been performed; however, there is no evidence to suggest carcinogenic potential.

Gamma glutamyl transpeptidase activity in rat urine is inhibited by various cephalosporins; however, the level of inhibition is less with cefuroxime. This may have significance in the interference in clinical laboratory tests in humans.

## **6. PHARMACEUTICAL PARTICULARS**

### **List of excipients**

None.

### **Incompatibilities**

None

### **Shelf life**

Unopened: 24 months

### **Special precautions for storage**

Store at a temperature below 30°C.

Protected from light. Reconstituted solution should be used immediately after preparation. Keep out of reach of children.

### **Nature and contents of container**

10ml Type III vial closed with 20mm Poly Bromo Rubber Stopper and 20mm Taxim blue flip off seal. Such 1 vial is packed in a mono carton along with 10ml sterile water for injection and pack insert.



**Special precautions for disposal and other handling**

None

**7. MARKETING AUTHORISATION HOLDER**

Sance Laboratories Private Limited.  
VI/51B, P.B. No: 2, Kozhuvanal-686573,  
Pala, Kottayam District, Kerala,  
**India.**

**8. MARKETING AUTHORISATION NUMBER(S)**

**TAN 21 HM 0091**

**9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORIZATION**

**March 19, 2021**

**10. DATE OF REVISION OF TEXT**

**March, 2021**